

Table 11. Estimates of daily wild rice consumption by the most abundant migratory birds encountered at Nett Lake, Minnesota in fall of 2007.

		Daily Wild Rice Consumption						
Common Name (reference)	Species	Per 1 bird in	Per 100 bird in	Per 1000 bird in	Per 5000 bird in	Per 10,000 bird in	Per 50,000 bird in	Per 100,000 bird in
		grams lbs	grams lbs	grams lbs	grams lbs	grams lbs	grams lbs	grams lbs
Ring-necked Duck (Hohman 1985) (Terrell and Wiser 1975)	<i>Aythya collaris</i>	27.90 0.06	2790.00 6.15	27900.00 61.52	139500.00 307.60	279000.00 615.20	1395000.00 3075.98	2790000.00 6151.95
Mallard (Jordan 1953)	<i>Anas platyrhynchos</i>	77.18 0.17	7718.00 17.02	77180.00 170.18	385900.00 850.91	771800.00 1701.82	3859000.00 8509.10	7718000.00 17018.19
Bluewinged Teal (Jordan 1953)	<i>Anas discors</i>	27.24 0.06	2724.00 6.01	27240.00 60.06	136200.00 300.32	272400.00 600.64	1362000.00 3003.21	2724000.00 6006.42
Canada Goose (Jordan 1953)	<i>Branta canadensis</i>	163.44 0.36	16344.00 36.04	163440.00 360.39	817200.00 1801.93	1634400.00 3603.85		
Redwinged Blackbird (Weatherhead et al. 1982) (Miller et al. 1950)	<i>Agelaius phoeniceus</i>	4.45 0.01	445.00 0.98	4450.00 9.81	22250.00 49.06	44500.00 98.12	222500.00 490.61	445000.00 981.23
Trumpeter Swan (Estimated from Jordan 1953)	<i>Cygnus buccinator</i>	550.00 1.21	55000.00 121.28	550000.00 1212.75				